



# The influence of maternal age, parity, blood group, gestational age and birth weight on fetomaternal haemorrhage in normal vaginal delivery

Studnickova M.<sup>1</sup>, Lubusky M.<sup>1,2</sup>, Simetka O.<sup>3</sup>, Prochazka M.<sup>1</sup>, Ordeltova M.<sup>4</sup>, Vomackova K.<sup>5</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, University Hospital, Olomouc, Czech Republic

<sup>2</sup>Department of Medical Genetics and Fetal Medicine, University Hospital, Olomouc, Czech Republic

<sup>3</sup>Department of Obstetrics and Gynecology, University Hospital, Ostrava, Czech Republic

<sup>4</sup>Department of Immunology, University Hospital, Olomouc, Czech Republic

<sup>5</sup>Department of Surgery I, University Hospital, Olomouc, Czech Republic

## AIM OF THE STUDY

Determine the influence of maternal age, parity, blood group, gestational age and birth weight on the volume of fetal erythrocytes, which enter the maternal circulation in normal vaginal delivery. Determining these parameters would enable improving the guidelines for RhD alloimmunization prophylaxis.

## WORKING HYPOTHESIS

A 10 µg dose of IgG anti-D administered intramuscularly should cover 0.5 ml of fetal RhD positive erythrocytes or 1 ml of whole blood. In the vast majority of pregnancies less than 2.5 ml of fetal erythrocytes (5 ml of whole blood, sufficient dose of IgG anti-D is 50 µg) enter the maternal circulation. Contrarily, only rarely does FMH exceed 5ml (10 ml of whole blood, sufficient dose of IgG anti-D is 100 µg) in normal vaginal delivery. Maternal age, parity, blood group, gestational age or birth weight does not influence the volume of fetomaternal hemorrhage in normal vaginal delivery.

## METHODS

A total of **3295** examinations were performed. The amount of fetal erythrocytes entering maternal circulation during uncomplicated spontaneous delivery of one fetus was determined by flow cytometry using the BDFACSCanto cytometer (Becton Dickonson International). Laboratory processing: Fetal Cell Count™ kit (Diagnosis of Fetomaternal Transfusion by flow cytometry), IQ Products, IQP-379. Calculation of total volume of fetal erythrocytes entering maternal circulation: Scientific Subcommittee of the Australian and New Zealand Society of Blood Transfusion. Guidelines for laboratory assessment of fetomaternal haemorrhage. 1st ed. Sydney: ANZSBT, 2002: 3-12.

## RESULTS

The average maternal age when FMH ≤ 1.7 ml (95 perc) was 29.4 years vs. 29.5 years when FMH > 1.7 ml, median 30 years in both groups, the difference was not statistically significant (p = 0.990).

The average gestational age when FMH ≤ 1.7 ml (95 perc) was 275.0 days vs. 276.5 days when FMH > 1.7 ml, median 278 days (39 weeks +5 days) vs. 277 days (39 weeks + 4 days), the difference was not statistically significant (p = 0.947).

The average birth weight when FMH ≤ 1.7 ml (95 perc) was 3309,6 g vs. 3328,9 g when FMH > 1.7 ml, median 3340 g vs. 3320 g, the difference was not statistically significant (p = 0.805).

FMH > 1.7 ml (5 perc) was present in 4.6% of primiparas (65/1418), in 4.8% of secundiparas (67/1307) and in 6.1% of multiparas (29/478), the difference was not statistically significant (p = 0.409).

FMH > 1.7 ml (5 perc) was present in 4.8% of maternal blood group 0 (53/1104), 5.1% A (67/1307), 3.6% B (22/617) and 7.1% AB (19/267), the difference was not statistically significant (p = 0.146).

FMH > 1.7 ml (5 perc) was present in 4.8% of maternal blood group RhD positive (120/2514) and 5.2% RhD negative (41/781), the difference was not statistically significant (p = 0.570).

The difference in maternal age, parity, blood group, gestational age and birth weight were also not statistically significant for fetomaternal hemorrhage FMH > 2.0 ml (2.5 perc), FMH > 2.5 ml (n = 34), FMH > 5 ml (n = 7).

## CONCLUSION

Maternal age, parity, blood group, gestational age and birth weight does not present a risk factor for excessive fetomaternal hemorrhage in normal vaginal delivery.

